

FIFRA Section 2(ee) Recommendation

For suppression of San Jose Scale (*Comstockaspis [Quadraspidotus] perniciosus*) in pome fruit, including apple, crabapple, pear, quince, loquat, and mayhaw.

For use in the State of NY only

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop: pome fruit, including apple, crabapple, pear, quince loquat, and mayhaw

Application Rate: Apply at a 1% v/v concentration beginning at crawler stage. Repeat on a 7-day interval during periods of scale infestation.

Crop-Specific Restrictions:

Do not apply when wind speed favors drift beyond the area intended for treatment.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

